

MILK FAT FREE FATTY ACIDS IN DEPENDENCE ON HEALTH OF DAIRY COWS

O. HANUŠ¹, M. KLIMEŠOVÁ¹, P. ROUBAL¹, E. SAMKOVÁ², D. FALTA³, M. ŠLACHTA², L. HASONŮVÁ²
 and I. NĚMEČKOVÁ¹

¹ *Dairy Research Institute Ltd., 160 00 Prague 6 – Vokovice, Czech Republic*

² *University of South Bohemia České Budějovice, Faculty of Agriculture, 370 05 České Budějovice, Czech Republic*

³ *Mendel's University in Brno, 613 00 Brno, Department of Animal Rearing and Breeding, Agronomical Faculty, Czech Republic*

Abstract

HANUŠ, O., M. KLIMEŠOVÁ, P. ROUBAL, E. SAMKOVÁ, D. FALTA, M. ŠLACHTA, L. HASONŮVÁ and I. NĚMEČKOVÁ, 2016. Milk fat free fatty acids in dependence on health of dairy cows. *Bulg. J. Agric. Sci.*, 22: 796–803

Free fatty acids (FFAs) in fat are important indicator of raw milk quality. The aim was to analyse the relationships and quantify effects of dairy cow mastitis on FFAs. Individual milk samples (IMs; n 307) came from Holstein and Czech Fleckvieh cows (67 : 33%). Sampling was from 10 to 270 days in milk for 1 year. Cows were fed in stables (n 5) by total mixed ration. Herd lactation milk yield varied from 7000 to 10 000 kg. Milking was performed twice a day using pipeline milking equipment and in milking parlours (20 : 80%). IMs were classified into 3 groups according to somatic cell count (SCC): A ≤ 150, healthy; B from 800 to 1500, subclinical mastitis; C > 1500 10³.ml⁻¹, subclinical and clinical mastitis. Lactose (L) decreased with growing intensity of mammary gland secretion disorder from 4.90 ± 0.278 (A) to 4.49 ± 0.44% (C; P < 0.001). The values increased in the same direction: content of FFAs from 0.65 ± 0.209 to 1.44 ± 0.379 mmol.100g⁻¹ (P < 0.001); milk energy quotients fat/protein (F/P) from 0.88 ± 0.386 to 1.23 ± 0.335 (P < 0.001) and fat/lactose (F/L) from 0.63 ± 0.266 to 1.98 ± 0.309 (P < 0.001). The mastitis state (SCC) deteriorated milk quality by increase of FFAs, 0.703 (P < 0.001; by 121.5%) and reduced L, -0.522 (P < 0.001). 49.4% of FFA variability could be explained by SCC variations. Increased SCC by 1000 10³.ml⁻¹ increased FFAs by 0.433 mmol.100g⁻¹. The L decrease due to worse mastitis state or cow energy malnutrition increased FFAs, -0.312 (P < 0.001). A risk increase of ketosis coefficients (F/P and F/L) deteriorated milk quality by increase of FFAs, 0.395 and 0.443 (P < 0.001) where 0.1 is equal to 0.05 and 0.067 mmol.100 g⁻¹. Values of FFAs can serve to control the health of dairy cows and quality of IMs in regular milk recording.

Key words: raw cow milk quality; mastitis; somatic cell count; lactose; ketosis quotient

Abbreviations: C – casein; CA – citric acid; CSN – Czech standard; F – fat; FFAs – free fatty acids; IMs – individual milk samples; IR – infrared; L – lactose monohydrate; MFP – milk freezing point depression; MI – milk indicator; MIR–FT – infrared spectrometry with Fourier's transformation; P – crude protein; PT – proficiency testing; SCC – somatic cell count; SNF – solids non fat; U – urea

Introduction

Current content of milk fat free fatty acids (FFAs) lies between 0.5 and 1.2 and maximal enabled is 1.3 for method by churning and 3.2 mmol.100 g⁻¹ for method by extraction

and titration according to CSN 57 0529 and CSN 57 0533. Increase of FFAs means row of negative impacts like metabolic problems of cows and enzymatic lipolysis (Figure 1). High FFAs cause an aggravation of milk technological properties (Vyletětlová et al., 1999, 2000 a, b) and deterioration of

*Corresponding author: hanus.oto@seznam.cz




Fig. 1. Lipolysis rise of fat in raw milk, content increase of free fatty acids (FFA), threat of milk and dairy product quality – factors and their combinations, related to animal and technology (modified according to Sjaunja (1984), O'Brian et al. (1998), Vyletřlová et al. (2000 a), Antonelli et al. (2002), Hanuš et al. (2004, 2008), Thomson et al. (2005), Deeth (2006); Wiking et al. (2006) and Mikulová (2011)

sensory milk properties. Therefore, it is important property for milk processing. Lipases can be thermoresistant and thus in this way to influence milk also after its heat treatment by dairy product degradation.

Wasteful milk handling as often pumping and ripple at manipulation (Sjaunja, 1984; Thomson et al., 2005; Hanuš et al., 2008) and its freezing on technology surfaces can induce a lipolysis. Poor hygiene of dairy cow stabling and milking as well as bad storage and treatment of raw milk can lead to propagation of undesirable psychrotrophic, thermoresistant and sporulating milk microflora (Vyletřlová et al., 1999; Cempřrková, 2002, 2007; Dankow et al., 2004; Hanuš et al., 2004; Deeth, 2006; Foltys and Kirchnerová, 2006; Hantis-Zacharov and Halpern, 2007; Cempřrková and Mikulová, 2009; Touřová et al., 2013). The mentioned facts can increase the lipolysis intensity (Figure 1).

Thus, results of FFAs are important at raw milk quality assessment for its processing on products. Despite of this fact these analyses are not carried out very often in bulk milk samples because of labour demandingness of original reference method in practice. Currently, after the start of MIR-FT (infrared spectrometry with Fourier's transformation) (Bijgaart van den, 2006; Hanuš et al., 2008, 2013 b) routine indirect analytical method, there is a new possibility to analyse larger sets of IMSs

as well. Thus, it will be possible to use the evaluated relationships between indicators of IMSs for advisory service to good milk quality and cow health at milk recording analyses.

The goal of this paper was to analyse and explain the basic relationships and quantify possible effects of dairy cow mastitis state and milk secretion disorders respectively on concentration of raw milk fat free fatty acids directly and indirectly in this way on quality and shelf life of milk products and milk food chain safety.

Materials and Methods

Animal conditions and individual milk samples

Individual milk samples (IMSs) were collected during whole year over all seasons ($n = 307$). These came from dairy cows of Holstein and Czech Fleckvieh breed in north Moravia in ratio 67 : 33%. Cows were sampled from early lactation to 270 days in milk. Daily milk yield of these animals varied from 10 to 34 kg according to lactation stage and curve. All animals were kept in stables for whole year without pasture and fed by total mixed ration (roughage as maize, alfalfa, clover and clover-grass silage and hay (specifically in dependence on herd, $n = 5$ with milk yield from 7000 to 10 000 kg per standard lactation) with supplements

of concentrates, mineral mixtures and vitamins according to current milk yield). Animals were kept in binding and free stables which were in ratio 20 : 80%. Dairy cows were fed and milked twice a day and milking was performed using pipeline milking equipment and in milking parlours (also 20 : 80%). The sampling was carried out using flow milkmeters (Tru-Test, New Zealand) and also dripping bottles in some milking parlours. IMSs were selected for experimental investigation and purposes using extremely marked somatic cell count ((SCC) ≤ 150 as healthy and $\geq 800 \cdot 10^3 \cdot \text{ml}^{-1}$ as suspect) and fat (F) > 1 and $< 7\%$ value because of possible highlighting of studied relationships (SCC classification) and avoidance of possible interference effects (F limitation). The samples were transported under refrigerator conditions ($< 8^\circ\text{C}$) in native state without preservation to laboratory and immediately analysed as follows.

Used milk indicators and relevant analytical methods

Infrared (IR) milk analyzer Lactoscope FTIR (Delta Instruments, The Netherlands) working in mid range of the IR spectrum with Fourier's transformation (MIR-FT) was used for determination of milk indicators (MI) such as F (fat), P (crude protein), C (casein), L (lactose monohydrate), SNF (solids non fat, all previous contents in %), U (urea concentration in $\text{mg} \cdot 100 \text{ ml}^{-1}$), CA (citric acid concentration in $\text{mmol} \cdot \text{l}^{-1}$), MFP (milk freezing point depression in $^\circ\text{C}$) and FFAs (milk fat content of free fatty acids in $\text{mmol} \cdot 100 \text{ g}^{-1}$). Milk SCC (somatic cell count in $10^3 \cdot \text{ml}^{-1}$) was investigated by the flow cytometry using Somacount FC (Bentley Instrument, USA). Both instruments were calibrated for various measurements of MI according to the results of relevant reference methods regularly with acceptable results. These instruments took part in the official proficiency testing (PT) of result reliability with good level. The MIR-FT measure-




Fig. 2. The result of Lactoscope (MIR-FT) calibration on measurement of FFAs ($\text{mmol} \cdot 100 \text{ g}^{-1}$) according to reference method results ($n = 6$; correlation coefficient 0.884, $P < 0.001$)

ment of FFAs was calibrated (Figure 2) (IDF 1991; Foss, 2001, 2004; Bijgaart van den, 2006) according to results of so called churning method (Koops et al., 1990; by titration using KOH solution) of determination of FFAs (CSN 57 0533 and 57 0529) with use of specific adjustment procedure (Hanus et al., 2008, 2013b). This instrument also took part in relevant PT on national level regularly with satisfactory evaluation.




Fig. 3. The relationship between somatic cell count (log SCC) and free fatty acids (FFAs; $\text{mmol} \cdot 100 \text{ g}^{-1}$) in chosen individual cow milk samples ($n = 307$; correlation index 0.703, $P < 0.001$)




Fig. 4. The relationship between somatic cell count (log SCC) and lactose (L; %) in chosen individual cow milk samples ($n = 307$; correlation coefficient -0.522 , $P < 0.001$)




Fig. 5. The relationship between lactose (L; %) and free fatty acids (FFAs; $\text{mmol} \cdot 100 \text{ g}^{-1}$) in chosen individual cow milk samples ($n = 307$; correlation coefficient -0.312 , $P < 0.001$)




Fig. 6. The relationship between fat/protein ratio (F/P) and free fatty acids (FFAs; $\text{mmol}\cdot 100\text{g}^{-1}$) in chosen individual cow milk samples ($n = 307$; correlation index 0.395, $P < 0.001$)




Fig. 7. The relationship between fat/lactose ratio (F/L) and free fatty acids (FFAs; $\text{mmol}\cdot 100\text{g}^{-1}$) in chosen individual cow milk samples ($n = 307$; correlation index 0.443, $P < 0.001$)

Table 1
Main statistic characteristics of indicators of individual milk samples ($n = 307$)

	x	xg	sd	v	min	max	m
SCC	2068		3577	173	5	43181	1221
log SCC	2.8666	736	0.744		0.699	4.6353	3.0867
F	3.85		1.298	33.7	1.03	6.97	3.81
P	3.51		0.431	12.3	2.58	4.7	3.45
L	4.68		0.39	8.3	2.88	5.35	4.73
SNF	8.81		0.478	5.4	7.56	10.13	8.81
C	2.79		0.369	13.2	1.93	3.93	2.73
U	38.37		9.734	25.4	7.11	68.77	39.12
CA	8.092		0.28	3.5	7.68	11.17	8.05
FFAs	1.2		0.499	41.7	0.12	3.14	1.18
F/P	1.11		0.38	34.3	0.29	2.09	1.12
F/L	0.84		0.322	38.4	0.21	1.97	0.82

x – arithmetic mean; xg – geometric mean; sd – standard deviation; v – variation coefficient (%); min – minimum; max – maximum; m – median; SCC – somatic cell count, $10^3\cdot\text{ml}^{-1}$; F – fat, %; P – crude protein, %; L – lactose monohydrate, %; STP – solids non fat, %; C – casein, %; U – urea, $\text{mg}\cdot 100\text{ml}^{-1}$; CA – citric acid, $\text{mmol}\cdot\text{l}^{-1}$; FFAs – free fatty acids, $\text{mmol}\cdot 100\text{g}^{-1}$; F/P – ratio fat/crude protein; F/L – ratio fat/lactose

SCC and FFAs data classification

Individual milk samples were classified and thus data about FFAs and other milk indicators were distributed into three groups according to SCC (as main indicator of milk secretion disorders and mammary gland health, in accordance) (Reneau et al., 1986; Pytlewski et al., 2012) values: A $\leq 150\ 10^3\cdot\text{ml}^{-1}$ as healthy mammary gland ($n = 94$); B from 800 to $1500\ 10^3\cdot\text{ml}^{-1}$ as problem with clear subclinical mastitis occurrence which can contribute to bulk milk as dairy plant delivery under practice conditions, it can influence of milk product quality ($n = 86$); C $> 1500\ 10^3\cdot\text{ml}^{-1}$ as trouble with subclinical and clinical mastitis occurrence and such milk is usually cancelled of dairy plant delivery and thus it does not affect milk product quality ($n = 127$).

Statistical procedures

Evaluation of FFAs MIR–FT calibration results was performed by linear regression in accordance with relevant system of statistical treatment (Figure 2). The main statistical parameters as means, standard deviations and variation coefficients were calculated (Table 1 and 2) for milk sample data sets and mean differences between SCC groups (A, B and C) of milk samples were tested by t–test (Table 3). Milk energy or ketosis quotients were calculated for milk sample results as ratio F/P and F/L. Logarithmic (\log_{10}) transformation of SCC data was performed because of absence of normal frequency distribution of these values (Reneau et al., 1986; Hanuš et al., 2008; Yilmaz and Koc, 2013) especially in individual milk samples where arithmetic mean is not suitable parameter for sets of SCC data and therefore in this way also geometric

means were used. Mutual relationships between milk indicators were evaluated by linear and nonlinear regression as well (Figure 3–7). The MS Excel (Microsoft, Redmond, USA) programme was used for this statistic evaluation.

Table 2

Main statistic characteristics of indicators of individual milk samples according to SCC groups

		n	x	xg	sd	v	min	max	m
A	SCC	94	82		37	45,1	5	149	82
	log SCC	94	1.8479	70	0.278		0.699	2.1732	1.9111
	F	94	3.05		1.256	41.2	1.03	6.25	2.82
	P	94	3.53		0.473	13.4	2.58	4.49	3.46
	L	94	4.9		0.278	5.7	3.69	5.35	4.94
	SNF	94	9.05		0.491	5.4	8.03	10.13	9
	C	94	2.8		0.412	14.7	1.93	3.64	2.74
	U	94	38.33		9.326	24.3	13.49	61.43	39.36
	CA	94	8.157		0.175	2.2	7.82	8.79	8.13
	FFAs	94	0.65		0.209	32.2	0.12	0.99	0.66
	F/P	94	0.88		0.386	44.1	0.29	1.93	0.81
	F/L	94	0.63		0.266	42.4	0.21	1.3	0.6
B	SCC	86	1112		206	18.5	808	1478	1109
	log SCC	86	3.0389	1094	0.081		2.9074	3.1697	3.0449
	F	86	4.03		1.131	28.1	1.44	6.97	3.88
	P	86	3.42		0.397	11.6	2.63	4.7	3.36
	L	86	4.7		0.261	5.5	3.78	5.25	4.75
	SNF	86	8.74		0.407	4.7	7.67	10.03	8.72
	C	86	2.72		0.336	12.3	2.05	3.93	2.68
	U	86	37.61		10.853	28.9	7.11	67.09	38.81
	CA	86	8.021		0.118	1.5	7.8	8.39	8.01
	FFAs	86	1.44		0.399	27.7	1	3.14	1.32
	F/P	86	1.18		0.322	27.2	0.45	2.09	1.18
	F/L	86	0.86		0.274	31.7	0.32	1.84	0.82
C	SCC	127	4186		4792	114.5	1508	43 181	2737
	log SCC	127	3.504	3192	0.279		3.1784	4.6353	3.4373
	F	127	4.32		1.159	26.8	1.58	6.82	4.26
	P	127	3.55		0.416	11.7	2.86	4.61	3.5
	L	127	4.49		0.44	9.8	2.88	5.19	4.6
	SNF	127	8.69		0.452	5.2	7.56	9.75	8.67
	C	127	2.83		0.352	12.5	2.24	3.77	2.77
	U	127	38.91		9.261	23.8	12.65	68.77	38.93
	CA	127	8.093		0.39	4.8	7.68	11.17	8.04
	FFAs	127	1.44		0.379	26.4	1	3.07	1.35
	F/P	127	1.23		0.335	27.3	0.39	2.01	1.19
	F/L	127	0.98		0.309	31.6	0.35	1.97	0.92

n – number of cases; A, $SCC \leq 150 \cdot 10^3 \cdot ml^{-1}$, healthy mammary gland; B, SCC from 800 to $1500 \cdot 10^3 \cdot ml^{-1}$, marked subclinical mastitis; C, $SCC > 1500 \cdot 10^3 \cdot ml^{-1}$, subclinical and clinical mastitis occurrence

Results and Discussion

The basic statistic results of data set (Table 1) about effect of SCC (sample groups A, B and C) on chosen milk

Table 3
Significance test of mean differences between SCC groups of individual milk samples

	A – B		A – C		B – C	
	t	sig	t	sig	t	sig
SCC	47.5	***	8.26	***	5.92	***
log SCC	38.03	***	43.51	***	14.99	***
F	5.44	***	7.73	***	1.81	ns
P	1.68	ns	0.38	ns	2.33	*
L	4.89	***	7.84	***	3.93	***
SNF	4.49	***	5.61	***	0.9	ns
C	1.44	ns	0.43	ns	2.14	*
U	0.48	ns	0.46	ns	0.94	ns
CA	6.01	***	1.46	ns	1.67	ns
FFAs	16.7	***	18.06	***	0.07	ns
F/P	5.74	***	7.14	***	0.9	ns
F/L	5.87	***	8.78	***	2.7	**

A – B, A – C and B – C – differences between SCC groups; t – t-test criterion; sig – difference significance; ns – insignificant, $P > 0.05$; * – significant, $P \leq 0.05$; ** – $P \leq 0.01$; *** – $P \leq 0.001$

indicators and especially on FFAs are shown in Table 2. In A group (healthy dairy cows), the largest variability of milk indicators was, in exception of SCC, in FFAs, fat, F/L and F/P (from 32.2 to 44.1%) and the lowest than in CA, SNF and L (from 2.2 to 5.7%) (Table 2). Statistic significance of mutual differences between milk indicator means of A, B and C group is shown in Tab. 3. Geometric means of SCC for A, B and C were 70, 1094 and 3192 $10^3 \cdot \text{ml}^{-1}$ (Table 2) and these were mutually different ($P < 0.001$; Table 3). This significance was logical in dependence on selection of IMSs. The significant differences (Table 3) at least between two groups were noted for most of observed milk indicators in dependence on mentioned distribution: L; SNF; casein; CA; FFAs; F/P; F/L. Only milk urea was not affected along groups due to carried out selection of samples (Table 2 and 3). Protein, casein and milk citric acid were influenced by mastitis state of dairy cows in less marked way.

Marked effects with clear trend caused by health SCC groups were noted in fat, SNF and especially in L, FFAs and milk ketosis quotients F/P and F/L (Table 2 and 3). Lactose decreased logically with growing intensity of mammary gland secretion disorder from 4.90 ± 0.278 (A) to $4.49 \pm 0.44\%$ (C; $P < 0.001$; Table 2 and 3). The following values increased also in the same direction with worsening health: content of FFAs from 0.65 ± 0.209 to 1.44 ± 0.399 (B) and 1.44 ± 0.379 $\text{mmol} \cdot 100 \text{ g}^{-1}$ ($P < 0.001$); milk energy quotients F/P from 0.88 ± 0.386 to 1.23 ± 0.335 ($P < 0.001$) and F/L from 0.63 ± 0.266 to 1.98 ± 0.309 ($P < 0.001$).

The chosen relationships between investigated indicators with respect to mammary gland health state and cow energy

metabolism milk indicators are shown in charts from Figures 3–7. The mastitis state (SCC) deteriorated milk quality markedly in terms of significant increase of FFAs with correlation index 0.703 ($P < 0.001$; Figure 3) and also reduced L with correlation coefficient -0.522 ($P < 0.001$; Figure 4) logically. The L decrease due to worsened mastitis state or possible cow energy malnutrition increased the FFAs, -0.312 ($P < 0.001$; Figure 5). On the contrary the risk increase both milk ketosis quotients (F/P and F/L) deteriorated milk quality by marked increase of FFAs with correlation indexes 0.395 and 0.443 (both $P < 0.001$) (Figure 6 and 7) similarly.

About possible mastitis (SCC) effect on content of FFAs was speculated already previously according to some papers (Figure 1) (Sjaunja, 1984; O'Brian et al., 1998; Vyletřlová et al., 2000a; Antonelli et al., 2002; Hanuš et al., 2004, 2008; Thomson et al., 2005; Wiking et al., 2006; Mikulová, 2011). However, the direct dependence with factor quantification has not been mentioned. Here, the experiment about quantification of this effect (Table 2) showed increase of FFAs by 121.5% as consequence of effects of subclinical (B) or subclinical and clinical mastitis (C). Technologically, it could be a part of spontaneous lipolysis (Toušová et al., 2013) but from physiological point of view this is induced lipolysis. 49,4% of FFA variability (Figure 3) could be explained by SCC variations. Increased SCC (due to mastitis) by 1000 $10^3 \cdot \text{ml}^{-1}$ increased FFAs by $0.433 \text{ mmol} \cdot 100 \text{ g}^{-1}$ (calculated from linear regression). Thomson et al. (2005) did not shown so close relationships in bulk milk samples although their relationships had the same trend in comparison to our findings. The carried out animal classification into groups (A,

B and C) with ambition to emphasize the effect of SCC on differences in milk quality (FFA) could be one of the reasons why here obtained closeness of relationships – Figure 3 (for SCC) and 5 (for L); 0.703 and -0.312 , $P < 0.001$) was markedly higher than in previous papers (for SCC 0.07 and for L -0.14 , $P < 0.05$ and $P < 0.01$) (Hanuš et al., 2008 and Thomson et al., 2005). The second reason is probably experiment with IMSs whereas other papers mentioned mostly bulk milk samples (Thomson et al., 2005; Hanuš et al., 2008; Mikulová, 2011). In these the impact can be soften down due to a diluting effect of healthy milk, it means without stress caused by subclinical production disorder occurrence. Lactose content which was reduced by secretion disorders (Figure 5) can explain 9.8% of variability in the occurrence of FFAs. With lactose decrease by 0.1% the FFA content grew up by $0.04 \text{ mmol} \cdot 100 \text{ g}^{-1}$.

As it has been defined (Geishhauser and Ziebell, 1995; Steen et al., 1996; Geishhauser et al., 1997; Duffield et al., 1997; Heuer et al., 2000; Siebert and Pallauf, 2010; van Knegsel et al., 2010; Hanuš et al., 2011, 2013a, c; van der Drift et al., 2012; Manzenreiter et al., 2013), milk F/P ratio is suitable indicator of dairy cow energy balance and its high value (for instance > 1.3) is indicator of suspicion from subclinical ketosis occurrence. Correlation between F/P and FFAs are not known for comparison purposes. Here, 39.5% of FFA variability (Figure 6) is explainable by variations in F/P ratio. Also milk F/L ratio is good indicator of dairy cow energy balance (Steen et al., 1996; Reist et al., 2002; Hanuš et al., 2013a, c; Manzenreiter et al., 2013). Also its higher value (for example > 0.9) is indicator of suspicion from subclinical ketosis occurrence. Correlation between F/L ratio and FFAs are not known for comparison purposes as well. Here, 19.6% of variability in FFAs (Figure 7) is explainable by variations in values of F/L. Therefore, the increase of F/P and F/L values (in linear interpretation) by 0.1 in consequence of nutritional energy problems of dairy cows is equal to the growth of FFAs by 0.05 and $0.067 \text{ mmol} \cdot 100 \text{ g}^{-1}$.

Conclusion

Significant relationships of SCC and both milk ketosis quotients (F/P and F/L) to FFAs show on diagnostical ability of FFA content not only regarding milk secretion disorders but also dairy cow energy troubles. Values of FFAs in IMSs (in regular milk recording) can serve to control the health of dairy cows in terms of suspicion from secretion or energy balance disorder occurrence or raw milk quality in consideration of quality and shelf-life of resulting milk products.

Acknowledgements

This research was supported by projects No. NAZV KUS QJ1210301, RO1415, NAZV QH 81210 and FA MENDELU TP 2/2013. Authors also thank to Mr. Jaroslav Kopecký and Mrs. Radoslava Jedelská for their support and technical cooperation.

References

- Antonelli, M. L., R. Curini, D. Scricciolo and G. Vinci, 2002. Determination of free fatty acids and lipase activity in milk: quality and storage markers. *Talanta*, **58**: 561–568.
- Bijgaart van den, H., 2006. New applications of mid-infra-red spectrometry for the analysis of milk and milk products. 2 Free fatty acids. *IDF Bulletin*, **406**: 22–28.
- Cempírková, R., 2002. Psychrotrophic vs. total bacterial counts in bulk milk samples. *Vet. Med.–Czech*, **47** (8): 227–233.
- Cempírková, R., 2007. Contamination of cow's raw milk by psychrotrophic and mesophilic microflora in relation to selected factor. *Czech J. Anim. Sci.*, **52** (11): 387–393.
- Cempírková, R. and M. Mikulová, 2009. Incidence of psychrotrophic lipolytic bacteria in cow's raw milk. *Czech J. Anim. Sci.*, **54** (2): 65–73.
- CSN 57 0529, 1993. Raw Cow Milk for Dairy Factory Treatment and Processing, ČNI, Praha.
- CSN 57 0533, 1997. Milk – Determination of Matter Content of Free Fatty Acids, ČNI, Prague.
- Dankow, R., J. Wojtowski and R. D. Fahr, 2004. Hygienic quality of raw milk in relation to methods of production and storage. *Med. Wet.*, **60**: 46–49.
- Deeth, H. C., 2006. Lipoprotein lipase and lipolysis in milk. *Inter. Dairy J.*, **16**: 555–562.
- Drift van der, S. G. K., R. Jorritsma, J. T. Schonewille, H. M. Knijn and J. A. Stegeman, 2012. Routine detection of hyperketonemia in dairy cows using Fourier transform infrared spectroscopy analysis of β -hydroxybutyrate and acetone in milk in combination with test-day information. *J. Dairy Sci.*, **95** (9): 4886–4898.
- Duffield, T. F., D. F. Kelton, K. E. Leslie, K. D. Lissimore and J. H. Lumsden, 1997. Use of test day milk fat and milk protein to detect subclinical ketosis in dairy cattle in Ontario. *Can. Vet. J.*, **38**: 713–718.
- Foltys, V. and K. Kirchnerová, 2006. Mesophilic and psychrotrophic aerobe sporulating microorganisms in raw cow's milk. *Centr. Eur. J. Biol.*, **1**: 545–560.
- FOSS, 2001. MilkoScan FT 6000; Free fatty acids measurement (FFA). *Spectrum Calibration*.
- FOSS, 2004. MilkoScan FT 120; Improved milk calibration. Analytical Application Note No. 128e.
- Geishhauser, T., K. E. Leslie, T. Duffield and V. Edge, 1997. Fat/protein ratio in first DHI tests milk as test for displaced abomasum in dairy cows. *Zbl. Vet. Med.*, **A 44** (5): 265–270.
- Geishhauser, T. and K. L. Ziebell, 1995. Fett/Eiweiss-Quotient in der Milch von Rinderherden mit Vorkommen von Labmagenverlagerungen. *Dtsch. Tierärztl. Wschr.*, **102**: 469–471.

- Hantš-Zacharov, E. and M. Halpern**, 2007. Culturable psychrotrophic bacteria communities in raw milk and their proteolytic and lipolytic traits. *Appl. Environ. Microbiol.*, **73**: 7162–7168.
- Hanuš, O., J. Frelich, M. Vyletětlová, P. Roubal, Z. Vorlíček and R. Jedelská**, 2004. Technologically difficult, pathogenic and food risky bacterial contamination of raw milk and other materials from dairy cow herds. *Czech J. Anim. Sci.*, **49** (11): 489–499.
- Hanuš, O., P. Roubal, E. Samková, D. Falta, G. Chládek and M. Vyletětlová–Klimešová**, 2013a. Cross validation of cut-off limit estimations for milk indicators of subclinical ketosis. *Acta Univ. Agric. et Silv. Mendel. Brun.*, **LXI** (6): 1661–1667.
- Hanuš, O., E. Samková, J. Říha, M. Vyletětlová–Klimešová and P. Roubal**, 2013b. Evaluation of development in indirect determination of milk fat free fatty acids in Czech Republic. *Acta Univ. Agric. et Silv. Mendel. Brun.*, **LXI** (6): 1669–1679.
- Hanuš, O., J. Vegrícht, J. Frelich, A. Macek, M. Bjelka, F. Louda and L. Janů**, 2008. Analyse of raw cow milk quality according to free fatty acids contents in the Czech Republic. *Czech J. Anim. Sci.*, **53** (1): 17–30.
- Hanuš, O., M. Vyletětlová–Klimešová, G. Chládek, P. Roubal and R. Seydlová**, 2013c. Metaanalysis of ketosis milk indicators in terms of their threshold estimation. *Acta Univ. Agric. et Silv. Mendel. Brun.*, **LXI** (6): 1681–1692.
- Hanuš, O., T. Yong, J. Kučera, V. Genčurová, A. Dufek, K. Hanušová and T. Kopec**, 2011. The predicative value and correlations of two milk indicators in monitoring energy metabolism of two breeds of dairy cows. *Sci. Agric. Boh.*, **42** (1): 1–11.
- Heuer, C., W. M. Van Straalen, Y. H. Schukken, A. Dirkszwaeger and J. P. T. M. Noordhuizen**, 2000. Prediction of energy balance in a high yielding dairy herd in early lactation: model development and precision. *Liv. Prod. Sci.*, **65**: 91–105.
- IDF**, 1991. Determination of free fatty acids in milk and milk products. *IDF Bulletin*, No. 256: Brussels.
- Knegsel van, A. T. M., S. G. A. Drift van der, M. Horneman, A. P. V. Roos de, B. Kemp and G. A. M. Graat**, 2010. Ketone body concentration in milk determined by Fourier transform infrared spectroscopy: Value for the detection of hyperketonemia in dairy cows. *J. Dairy Sci.*, **93**: 3065–3069.
- Koops, J., H. Klomp and H. Hemert van**, 1990. Rapid enzymatic assay of free fatty acids (lipolysis) in farm tank milk by a segmented continuous-flow method. Comparison of the results with those obtained by the BDI procedure. *Neth. Milk Dairy J.*, **44**: 3–19.
- Manzenreiter, H., B. Fürst-Waltl, C. Egger-Danner and W. Zollitsch**, 2013. Zur Eignung des Gehalts an Milchinhaltsstoffen als Ketoseindikator. 40 *Viehwirtschaft. Fachtag.*, 9–19.
- Mikulová, M.**, 2011. Content of free fatty acids lipolytic bacteria and somatic cells in relation to milking technology. *J. Agrobiol.*, **28** (1): 49–54.
- O'Brian, B., E. O'Callaghan and P. Dillon**, 1998. Effect of milking machine systems and components on free fatty acid levels in milk. *J. Dairy Res.*, **65**: 335–339.
- Pytlewski, J., I. Antkowiak, M. Adamski, J. Kučera and R. Skrzypek**, 2012. Factors associated with hygienic quality of bulk tank milk produced in central Poland. *Ann. Anim. Sci.*, **12** (2): 227–235.
- Reist, M., D. von Euw, K. Tschuemperlin, L. Leuenberger, Y. Chilliard, H. M. Hammon, C. Morel, C. Philipona, Y. Zbinden, N. Kuenzi and J. W. Blum**, 2002. Estimation of energy balance at the individual and herd level using blood and milk traits in high-yielding dairy cows. *J. Dairy Sci.*, **85**: 3314–3327.
- Reneau, J. K.**, 1986. Effective use of dairy herd improvement somatic cell counts in mastitis control. *J. Dairy Sci.*, **69** (6): 1708–1720.
- Siebert, F. and J. Pallauf**, 2010. Analyse von Ergebnissen der Milchleistungsprüfung in Hessen im Hinblick auf ein Ketosorisiko. *Züchtungskunde*, **82**: 112–122.
- Sjaunja, L. O.**, 1984. Studies on milk analysis of individual cow milk samples. III. The effect of different treatments on infrared analyses. *Acta Agric. Scand.*, **34**: 273–285.
- Steen, A., O. Osteras and H. Gronstol**, 1996. Evaluation of additional acetone and urea analyses, and of the fat–lactose–quotient in cow milk samples in the herd recording system in Norway. *J. Vet. Med.*, **A 43**: 181–191.
- Thomson, N. A., W. M. Woolford and A. P. J. Copeman**, 2005. Milk harvesting and cow factors influencing seasonal variation in the levels of free fatty acids in milk from Waikato dairy herds. *New Zeal. J. Agr. Res.*, **48**: 11–21.
- Toušová, R., L. Stádník and J. Ducháček**, 2013. Effects of season and time of milking on spontaneous and induced lipolysis in bovine milk fat. *Czech J. Food Sci.*, **31** (1): 20–26.
- Vyletětlová, M., P. Benda, O. Hanuš and P. Kopunecz**, 1999. Determination of total psychrotrophic microorganisms in bulk milk samples and their relationship to total count of mikroorganisms. *Czech J. Food Sci.*, **17** (6): 216–222. (Cz).
- Vyletětlová, M., J. Ficnar and O. Hanuš**, 2000 a. Effects of lipolytic enzymes *Pseudomonas fluorescens* on liberation of fatty acids from milk fat. *Czech J. Food Sci.*, **18** (5): 175–182. (Cz)
- Vyletětlová, M., O. Hanuš, E. Urbanová and P. Kopunecz**, 2000 b. The occurrence and identification of psychrotrophic bacteria with proteolytic and lipolytic activity in bulk milk samples at storage in primary production conditions. *Czech J. Anim. Sci.*, **45**: 373–383. (Cz)
- Wiking, L., J. H. Nielsen, A. K. Bavius, A. Edvardsson and K. Svennersten-Sjaunja**, 2006. Impact of milking frequencies on the level of free fatty acids in milk, fat globule size, and fatty acid composition. *J. Dairy Sci.*, **89**: 1004–1009.
- Yilmaz, H. and A. Koc**, 2013. A research on milk yield, persistency, milk constituents and somatic cell count of Red Holstein cows reared under mediterranean climatic conditions. *Bulg. J. Agric. Sci.*, **19** (6): 1401–1407.